

We claim:

- A process for the preparation of dimethylcumenes comprising alkylating a substrate comprising of one or more xylene isomers with an alkylating agent in the presence of a solid acid zeolite catalyst selected from ultrastable zeolite Y (Si/Al = 5 to 50) and Beta (Si/Al = 10 120), and separating the products formed in vapour phase.
- 2. A process as claimed in claim 1 wherein said substrate and alkylating agent are contacted with said solid acid zeolite catalyst at a temperature in the range of 80 250°C and for a period of at least 1 hour.
- 3. A process as claimed in claim 1 wherein the product is separated from the vapour phase by condensation at a temperature in the range of 0 3°C.
- 4. A process as claimed in claim 1 wherein the substrate is selected from o-xylene, m-xylene, p-xylene and any mixture thereof.
- 5. A process as claimed in claim 1 wherein the Si/Al ratio in said catalyst is between 5 to 20.
- 6. A process as claimed in claim 1 wherein the alkylating agent is selected from propylene and propyl alcohols such as isopropanol and n-propanol.
- 7. A process as claimed in claim 2 wherein the temperature of the reaction is in the range of 100 200°C.
- 8. A process as claimed in claim 2 wherein the temperature of the reaction is in the range of 120 180°C.
- 9. A process as claimed in claim 1 wherein the molar ratio of xylene substrate to the alkylating agent in the feed is in the range of from 1:2 to 20:1.
- 10. A process as claimed in claim 9 wherein the molar ratio of xylene substrate to the alkylating agent is in the range of 1:1 to 10:1.
- 11. A process as claimed in claim 9 wherein the molar ratio of xylene substrate to the alkylating agent is in the range of 1:2 to 5:1.
- 12. A process as claimed in claim 1 wherein the weight hourly space velocity (WHSV) of the feed is in the range of 0.5 to 30 h⁻¹.
- 13. A process as claimed in claim 12 wherein the weight hourly space velocity (WHSV) of the feed is in the range of 1 to 20h⁻¹.
- 14. A process as claimed in claim 12 wherein the weight hourly space velocity (WHSV) of the feed is in the range of 2 to 10h⁻¹.
- 15. A process as claimed in claim 1 wherein the alkylation reaction is carried out in a fixed bed reactor or a batch reactor.





- 16. A process as claimed in claim 1 wherein p-xylene is alkylated using isopropanol in the presence of zeolite beta catalyst.
- 17. A process as claimed in claim 1 wherein m-xylene and o-xylene are alkylated using isopropanol as the alkylating agent in the presence of ultrastable zeolite Y (USY) as catalyst.
 - 18. A process as claimed in claim 1 wherein a mixture of *p*-xylene and isopropyl alcohol in a molar ratio of 4:1 is reacted in a fixed bed reactor in the presence of ultrastable zeolite Y catalyst.